

Supplemental Table 1. Immunoglobulin levels.

	Initial	Post-RFCM	Post-Rituximabmaintenance	P*
IgA < 0.7 mg/dL	49.1% (28/57)	42.4% (25/59)	58.8% (30/51)	0.13
IgG < 6.5 mg/dL	36.8% (21/57)	33.9% (20/59)	49% (25/51)	<0.001
IgM < 0.4 mg/dL	62.5% (35/56)	76.3% (45/59)	88.2% (45/51)	<0.001

**T test paired samples comparing initial values and final values*

Supplemental Table 2. Pharmacokinetic parameters obtained for each patient.

	Patient 21(PR) ^{&}	Patient 48 (CR) [§]	Patient 19 (CR)	Patient 23 (CR)	Patient 34 (CR)	Patient 8 (CR)	Mean (SD)
Dose (mg)	589	750	664	645	750	810	701.3 (82.1)
AUC (μg $\cdot\text{day/L/1.73m}^2$)	1184.4	2328.2	3577.4	3158.1	3131.5	1983.6	2675.9 (948.4)
Cl (mL/day/1.73m^2)	603.8	278.7	180.0	204.2	188.5	258.2	285.6 (160.8)
Vc (L/1.73m^2)	3037.6	3354.2	3035.1	5705.9	4063.9	3889.0	3847.6 (1006.1)
Cmax ($\mu\text{g/mL}$)	219.3	193.4	212.2	113.0	159.5	163.7	176.9 (39.7)
Cmin ($\mu\text{g/mL}$)	1.0	2.4	7.2	10.1	11.1	5.4	6.2 (4.1)
T1/2 (days)	3.4	8.3	11.7	19.4	14.9	10.4	11.4 (5.5)
T1/2 α (days)	3.3	0.7	1.1	9.5	1.9	1.0	2.9 (3.4)
T1/2 β (days)	3.4	17.1	23.3	50.5	31.9	21.6	24.6 (15.8)
Vss (L/1.73m^2)	2960.5	6592.7	5779.7	10360.0	8557.3	7635.6	6981.0 (2533.3)
MRT (days)	4.9	23.7	32.2	50.7	42.9	29.6	30.7 (15.9)

[&]Partial responder, [§]Complete responder