


Supplemental Table: Population Characteristics

	Patients Discharged Without Anticoagulation (N=163)
Age, year (Median, IQR)	58 (44-67)
Sex: Female	85 (52.2%)
Race: White African American Asian Unknown	57 (36.5%) 62 (39.7%) 6 (3.9%) 31 (19.9%)
Comorbidities Hypertension Diabetes Heart Disease Chronic Respiratory Disease Cancer Chronic Kidney Disease Liver Disease Pregnancy	77 (52.8%) 51 (31.3%) 20 (12.2%) 37 (22.4%) 23 (14.1%) 16 (9.8%) 9 (5.5%) 15 (9.2%)
Hospitalization and Discharge Details	
D-Dimer on Admission (Median, IQR)	856 (528-1428)
Required intensive care during index admission	42 (25.8%)
Length of Stay, days (Median, IQR)	6 (3-12)
Anticoagulation During Hospitalization* :	
No Anticoagulation	7 (4.3%)
Prophylactic dose	152 (93.6%)
Intermediate-dose	22 (13.5%)
Therapeutic	18 (11.1%)
Duration of Follow Up recorded (Median, IQR)	30 (17-46)
Type of Last Contact Recorded:	
Telehealth	115 (69.7%)
In person Visit	36 (22%)
Readmission	12 (7.3%)

* Some patients received more than one anticoagulant dose during the course of hospitalization.



Supplementary Figure: Consort Diagram